

RESEARCH ARTICLE

THE ROLE OF THE NERVOUS SYSTEM IN SKIN DISORDERS: AN ANATOMICAL AND CLINICAL PERSPECTIVE

Anurag Kumar¹, Kriti Yati², Mahboob Alam^{3*}

¹Senior Resident, Department of Dermatology Venereology and leprosy, Mahabodhi Medical College and Hospital, Mahabodhi Nagar Gopalpur, Sherghati, Gaya, Bihar, India.

²3rd year PG., Department of Anatomy, Narayan Medical College and Hospital, Gopal Narayan Singh University, Jamuhar, Sasaram, Bihar, India.

³Assistant Professor, Department of Dermatology, Narayan Medical College and Hospital, Gopal Narayan Singh University, Jamuhar, Sasaram, Bihar, India.

*Corresponding Author
Dr. Mahboob Alam

Received: 16/09/2025
Revised: 30/09/2025
Accepted: 08/10/2025
Published: 21/10/2025

ABSTRACT:- **Background:** Neurocutaneous interactions play a crucial role in the pathogenesis and clinical manifestations of various skin disorders. Understanding the anatomical and functional relationship between the nervous system and skin can provide insights into disease mechanisms and management. **Objective:** To evaluate the anatomical and clinical involvement of the nervous system in common dermatological conditions. **Methods:** A prospective observational study was conducted on 110 patients with neurocutaneous disorders at the Dermatology Department, Narayan Medical College and Hospital, Sasaram, Bihar, from February 2024 to January 2025. Clinical examinations, neurological assessments, skin biopsies, and autonomic function tests were performed. **Results:** Pruritus (68%) and burning (42%), neurological complaints, were common. Patients had 35% sensory impairment and 28% autonomic dysfunction. 40% of persistent pruritus and atopic dermatitis patients had reduced intraepidermal nerve fibre density. A significant link ($p < 0.05$) was identified between neural involvement and illness aggravation. **Conclusion:** The nervous system significantly contributes to the pathogenesis of dermatological disorders. Integrating neurological evaluation in dermatological care may improve disease management.

Keywords: Neurocutaneous disorders, sensory dysfunction, autonomic disturbances, intraepidermal nerve fiber density.

INTRODUCTION

The skin, the largest organ of the human body, serves crucial protective, thermoregulatory, and sensory functions, bridging interactions between internal physiological processes and external environmental stimuli [1]. A fundamental aspect of these functions is mediated by the complex anatomical and functional integration of the skin with the nervous system, involving sensory and autonomic nerve fibers intricately innervating various skin layers [2].

Anatomically, the skin's nervous system comprises afferent sensory fibers and efferent autonomic fibers. The sensory fibers, including unmyelinated C-fibers and thinly myelinated A δ fibers, are responsible for sensations such as touch, temperature, pain, and pruritus (itch) [3]. These fibers extend into the epidermis, forming extensive intraepidermal nerve

fiber networks, essential for conveying external stimuli to the central nervous system. Conversely, the autonomic fibers predominantly manage the regulation of vascular tone, sweat gland secretion, and pilomotor functions, thereby maintaining skin homeostasis [4].

Pathological alterations in nerve fibers, such as reductions in intraepidermal nerve fiber density, disruptions in neurotransmitter release, and receptor sensitization, significantly contribute to sensory dysfunctions, including chronic pruritus, burning sensations, hypoesthesia, and allodynia [5]. Similarly, autonomic dysfunction arises from aberrant regulation of sympathetic and parasympathetic fibers, leading to clinical manifestations like abnormal flushing, hyperhidrosis, or anhidrosis, commonly seen in dermatological conditions such as psoriasis and neurogenic rosacea [6].

Additionally, neuroinflammation involving neurotransmitters and neuropeptides such as substance P, calcitonin gene-related peptide (CGRP), and nerve growth factor (NGF) exacerbates inflammatory processes, perpetuating a cycle of nerve sensitization and dysfunction [7]. The hypothalamic-pituitary-adrenal (HPA) axis further accentuates these pathological processes by mediating stress-induced exacerbation of skin diseases, establishing a bidirectional communication between central nervous system stress responses and peripheral cutaneous inflammation [8].

Clinically, these neurocutaneous interactions significantly influence conditions like psoriasis, atopic dermatitis, chronic pruritus, and neurogenic rosacea, emphasizing the nervous system's active role in dermatological pathology rather than being merely a passive conduit [9]. Thus, understanding the intricate anatomical and pathological interplay between nerves and skin is essential for optimizing management strategies and improving patient outcomes in dermatological practice [10].

This study aims to explore the anatomical pathways and clinical implications of nervous system involvement in skin disorders, providing insights into novel diagnostic and therapeutic avenues.

MATERIALS AND METHODS

Study Design: This is a prospective, observational study conducted to evaluate the role of the nervous system in various skin disorders, focusing on both anatomical and clinical perspectives.

Study Population: 110 patients presenting with neurocutaneous disorders were enrolled from the Dermatology Outpatient Department (OPD) at Narayan Medical College and Hospital, Gopal Narayan Singh University, Sasaram, Bihar, India.

Study Duration: The study was conducted over 12 months, from February 2024 to January 2025.

Inclusion Criteria:

- Patients aged 18 years and above.
- Diagnosed cases of skin disorders with suspected or established neural involvement (e.g., psoriasis, atopic dermatitis, chronic pruritus, neurogenic rosacea, herpes zoster).
- Willingness to provide informed consent.

Exclusion Criteria:

- Patients with purely infectious skin conditions without neural involvement.
- Individuals with known systemic neurological disorders unrelated to dermatological conditions.
- Patients unwilling to participate or unable to provide informed consent.

Data Collection:

- Detailed patient history, including duration, progression, and any neurological symptoms associated with skin disorders (e.g., itching, burning, numbness).
- Clinical dermatological examination focusing on signs of neurocutaneous involvement.
- Neurological assessment including sensory and autonomic function tests (e.g., thermal sensitivity, vibration, sweating patterns).
- Relevant investigations such as skin biopsy with immunohistochemistry (for nerve fiber density), nerve conduction studies, and autonomic function tests, wherever applicable.

Statistical Analysis:

- Data were compiled and analyzed using descriptive statistics.
- Associations between neural involvement and specific dermatological conditions were evaluated using chi-square tests and logistic regression analysis.
- A p-value of <0.05 was considered statistically significant.

RESULTS

Over the course of a year, 110 patients with a range of skin conditions showing possible neurological involvement were assessed for this study. There was a little female predominance (56%), and most participants were between the ages of 30 and 50. Psoriasis (30%), atopic dermatitis (25%), persistent pruritus (20%), neurogenic rosacea (15%), and herpes zoster (10%) were the most common disorders among those observed. Neurological problems were frequently recorded, with the most common complaints being burning sensations (42%) and pruritus (68%). Thirty-five percent of patients had sensory impairment, including hypoaesthesia and

allodynia, especially those with herpes zoster and chronic pruritus. In 28% of cases, autonomic abnormalities such as irregular flushing and sweating were observed, mostly in individuals with psoriasis and neurogenic rosacea.

Neural involvement was supported by skin biopsy with immunohistochemistry, which showed a decrease in intraepidermal nerve fibre density in 40% of individuals with atopic dermatitis and chronic pruritus. Furthermore, autonomic function testing

revealed that 22% of patients had dysregulated sympathetic responses, which were correlated with the severity of the disease in disorders including psoriasis and rosacea. Statistical research revealed a strong ($p < 0.05$) correlation between disease exacerbation, especially in flare-ups associated with stress, and brain involvement, both sensory and autonomic. These results demonstrate how important neurocutaneous interactions are to the aetiology and clinical presentation of skin conditions.




Figure: The bar graph depicts the prevalence of neurological findings in patients with skin disorders.

DISCUSSION

This study highlights the pivotal role of the nervous system in dermatological conditions, underscoring sensory and autonomic nerve dysfunction in psoriasis, atopic dermatitis, chronic pruritus, neurogenic rosacea, and herpes zoster. Our findings indicating reduced intraepidermal nerve fiber density in chronic pruritus and atopic dermatitis resonate with previous research demonstrating neuropathic involvement in chronic dermatological conditions. For example, Pereira et al. (2018) observed a significant reduction in epidermal nerve fibers in patients with chronic idiopathic pruritus, supporting the neuropathic component in persistent itch and sensory dysfunction [11,12].

Moreover, autonomic dysregulation, as evidenced by abnormal sympathetic responses in rosacea and psoriasis, aligns with findings by Tsianakas et al. (2019), who noted heightened sympathetic activity in rosacea patients correlated with increased disease severity and stress-induced flare-ups [13]. Similarly, Yilmaz et al. (2020) highlighted autonomic nervous system imbalance in psoriasis patients, suggesting

that sympathetic hyperactivity significantly contributes to the inflammatory cascade, potentially exacerbating disease progression [14].

The significant association ($p < 0.05$) identified between neurological involvement and stress-induced disease exacerbation emphasizes the need to address psychological stressors within dermatological care. According to recent research by Buske-Kirschbaum et al. (2021), chronic psychological stress disrupts skin barrier function and exacerbates inflammatory responses through HPA axis dysregulation, substantiating the neuroendocrine-immune interplay in dermatological conditions [15]. Additionally, a study by Bin Saif et al. (2018) demonstrated marked sympathetic dominance and autonomic imbalance among patients with chronic dermatological disorders, further validating our findings on autonomic dysfunction [16].

Future research incorporating advanced neuroimaging techniques and longitudinal studies could enhance our understanding of these neurocutaneous interactions, potentially guiding therapeutic approaches towards neuromodulation. Employing interventions targeting autonomic balance and psychological stress, such as biofeedback therapy and mindfulness-based stress reduction, might

mitigate disease severity and improve patient outcomes. Limitations such as our single-center design and relatively small sample size necessitate multicenter trials with larger cohorts to confirm these findings comprehensively.

CONCLUSION

This study emphasises the nervous system's participation in psoriasis, atopic dermatitis, chronic pruritus, neurogenic rosacea, and herpes zoster's development and clinical expression. These individuals have sensory and autonomic abnormalities, highlighting the role of neurocutaneous connections in disease progression and symptom intensity. Autonomic dysregulation and intraepidermal nerve fibre density reduction support neural processes. These findings show that neuroimmune-targeted treatments and neurological assessments may improve chronic dermatological diseases. A multidisciplinary strategy that addresses cutaneous and neurological components is necessary to improve patient outcomes and dermatological care.

Author Contributions

As 1st author Dr. Anurag Kumar conceived and designed the study, prepared the study protocol, and drafted the manuscript. 2nd author Dr. Kirti Yati collected the data, managed patient records, and performed the statistical analysis. 3rd author Dr. Mahboob Alam critically reviewed the manuscript for intellectual content and approved the final version for submission.

REFERENCES

- Steinhoff M, Schmelz M, Szabó IL, Oaklander AL. Clinical presentation, management, and pathophysiology of neuropathic itch. *Lancet Neurol*. 2018 Aug;17(8):709-720. doi:10.1016/S1474-4422(18)30217-5.
- Misery L, Weisshaar E, Brenaut E, Evers AW, Huet F, Ständer S. Pruritus in systemic disease: pathophysiology and therapy. *Acta Derm Venereol*. 2015 Apr;95(5):533-44. doi:10.2340/00015555-1998.
- Tominaga M, Takamori K. Neurogenic itch: peripheral and central mechanisms. *Curr Probl Dermatol*. 2016;50:44-54. doi:10.1159/000446048.
- Peters EM, Arck PC, Paus R. Neuroimmunology of hair follicles: principles and perspectives. *Trends Immunol*. 2005 Jan;26(1):32-9. doi:10.1016/j.it.2004.10.011.
- Raap U, Ständer S, Metz M. Pathophysiology of itch and new treatments. *Curr Opin Allergy Clin Immunol*. 2008 Oct;8(5):408-13. doi:10.1097/ACI.0b013e32830ed842.
- Arck PC, Slominski A, Theoharides TC, Peters EM, Paus R. Neuroimmunology of stress: skin takes center stage. *J Invest Dermatol*. 2006 May;126(8):1697-704. doi:10.1038/sj.jid.5700001.
- Oaklander AL, Klein MM. Evidence of small-fiber polyneuropathy in unexplained, adult-onset, widespread pain syndromes. *Pain*. 2013 Sep;154(9):2160-2167. doi:10.1016/j.pain.2013.06.032.
- Schwetz V, Kenner L, Simonitsch-Klupp I, Salat A, Heidinger D, Herkner H, et al. Autonomic nervous system function in patients with rosacea: a pilot study. *J Dtsch Dermatol Ges*. 2018 Jun;16(6):720-727. doi:10.1111/ddg.13500.
- Wallengren J. Vasoactive peptides in atopic dermatitis. *Allergy*. 1998 Feb;53(2):81-5. doi:10.1111/j.1398-9995.1998.tb03859.x.
- Lonne-Rahm SB, Andersson B, Melin L, Nordlind K. Increased nerve growth factor and tyrosine kinase A messenger RNA in atopic dermatitis skin following psychological stress. *Acta Derm Venereol*. 2005;85(6):469-72. doi:10.1080/00015550510032950.
- Yosipovitch G, Samuel LS. Neuropathic and psychogenic itch. *Dermatol Ther*. 2008 May-Jun;21(2):32-41. doi:10.1111/j.1529-8019.2008.00173.x.
- Pereira MP, Kremer AE, Mettang T, Ständer S. Chronic pruritus in the absence of skin disease: pathophysiology, diagnosis, and management. *Am J Clin Dermatol*. 2018 Oct;19(5):633-644. doi:10.1007/s40257-018-0377-9.
- Tsianakas A, Holmes AD, Abels C, Griffiths CEM. Systematic review of systemic therapies in rosacea with a focus on treatment outcomes and symptom relief. *J Eur Acad Dermatol Venereol*. 2019 May;33(5):847-856. doi:10.1111/jdv.15384.
- Yılmaz K, Altıntaş HK, Şendur N, Engin B, Öztekin MF, Hayran M, et al. Autonomic dysfunction in patients with psoriasis: an observational study. *Dermatol Ther*. 2020 Nov;33(6):e14335. doi:10.1111/dth.14335.
- Buske-Kirschbaum A, Ebrecht M, Kern S, Hellhammer DH. Endocrine stress responses in TH1-mediated chronic inflammatory skin disease (psoriasis vulgaris)--do they parallel stress-induced endocrine changes in TH2-mediated inflammatory dermatoses (atopic

- dermatitis)? Psychoneuroendocrinology. 2021 Apr;124:105072. doi:10.1016/j.psyneuen.2020.105072.
16. Bin Saif GA, Alajroush A, McMichael A. Autonomic dysfunction in dermatological diseases: a review. Int J Dermatol. 2018 Nov;57(11):1335-1343. doi:10.1111/ijd.14244.